

This article was downloaded by:[Uribe, Sergio]  
On: 6 February 2008  
Access Details: [subscription number 790434916]  
Publisher: Informa Healthcare  
Informa Ltd Registered in England and Wales Registered Number: 1072954  
Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Acta Odontologica Scandinavica

Publication details, including instructions for authors and subscription information:  
<http://www.informaworld.com/smpp/title~content=t713394069>

### "Risk periods" associated with the development of dental fluorosis in maxillary permanent central incisors: a meta-analysis

Asgeir Bårdsen

Online Publication Date: 07 February 1999

To cite this Article: Bårdsen, Asgeir (1999) "Risk periods" associated with the development of dental fluorosis in maxillary permanent central incisors: a meta-analysis', Acta Odontologica Scandinavica, 57:5, 247 - 256

To link to this article: DOI: 10.1080/000163599428652

URL: <http://dx.doi.org/10.1080/000163599428652>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article maybe used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

# “Risk periods” associated with the development of dental fluorosis in maxillary permanent central incisors: a meta-analysis

Asgeir Bårdsen

Department of Dental Research, Faculty of Dentistry, University of Bergen, Bergen, Norway

Bårdsen A. “Risk periods” associated with the development of dental fluorosis in maxillary permanent central incisors: a meta-analysis. *Acta Odontol Scand* 1999;57:247–256. Oslo. ISSN 0001-6357.

This systematic review of the dental literature (1966–98) concerns risk periods associated with dental fluorosis in the maxillary permanent central incisors. A literature search was organized through the MedLine and the ISI databases. In addition, one unpublished paper (in manuscript) was obtained, as well as one paper published before 1966. However, out of 143 catches, only 10 studies were included in this review. The main reason for exclusions was that the data presented did not meet the criteria given for the present meta-analysis. Among the included papers, 7 pertained to subjects whose exposure to fluoride started at different ages during the enamel formation (Group 1), and 3 were based on subjects who had been exposed from birth and then experienced an abrupt reduction in daily fluoride exposure at different ages during the amelogenesis period (Group 2). The meta-analysis for Group 1 found the odds ratio (OR) for dental fluorosis in children exposed to fluoride early in life (before 2 years of age) to be 7.24 (95% CI; 4.71–11.13) as compared to children exposed later in life (after 2 years of age). The meta-analysis for the studies in Group 2 found the overall OR to be 1.88 (95% CI; 1.35–2.61) for children who had a reduction in fluoride intake after 2 years of age, as compared to individuals who experienced reduction earlier (during the first 2 years). The studies from both groups were pooled and the duration of exposure to fluoride during the first 4 years of life was the independent variable. The meta-analysis now revealed an overall OR of 5.83 (95% CI; 2.83–11.94) for long periods of fluoride exposure (>2 out of the first 4 years) versus shorter periods of exposure (<2 out of the first 4 years of life) during the enamel formation in the maxillary central incisors. Based on the findings of the meta-analysis, no specific period of enamel formation is singled out as being the most critical for the development of dental fluorosis. The duration of fluoride exposure during the amelogenesis, rather than specific risk periods, would seem to explain the development of dental fluorosis in the maxillary permanent central incisor. □ *Dental fluorosis; fluoride exposure; meta-analysis; risk period; susceptibility*

Asgeir Bårdsen, Department of Dental Research, Faculty of Dentistry, University of Bergen, Årstadveien 17, N-5009 Bergen, Norway. Tel. +47 55 58 64 74, fax. +47 55 58 64 87, e-mail. asgeir.bardsen@odont.uib.no

Excessive fluoride intake can cause dental fluorosis in developing teeth (1–3). While the clinical features of dental fluorosis are well known, the exact role of fluoride in the pathogenesis and the period(s) of maximum risk are still matters of debate. Relatively few studies have addressed the problem, and findings are inconsistent. Since fluorotic changes in teeth cannot be cured, but may easily be prevented if the fluoride intake is adequately controlled during the period of tooth formation, more information on the timing of dental fluorosis is urgently needed.

The start and duration of the mineralizing period vary among the different groups of teeth, so the period of maximum susceptibility would be expected to vary accordingly. In order to simplify the assessment of risk periods, the present study is restricted to maxillary permanent central incisors, in which mineralization of the crown is normally completed by 4 years of age (4).

The ideal study regarding fluorosis risk periods in the incisors would be a randomized clinical trial, including groups of subjects exposed to fluoride only during the 1st year of life, alternatively only the 2nd, 3rd or 4th year of life. Ethical and practical considerations make it impossible to conduct such experiments in humans. Information concerning risk periods in humans can, consequently, only

be gathered retrospectively; from individuals who have been exposed to changes in the fluoride ingestion more or less by chance.

The literature concerning the timing of fluorosis includes at least two types of reports: (i) studies in individuals who have been introduced to fluoride at different ages during their first years of life (5–9) and (ii) studies in populations which, after having been exposed to fluoride (water fluoride) from birth, experience an abrupt reduction in daily fluoride intake at different periods during the amelogenesis (10–12).

Due to methodological differences in the studies (grouping, indices for scoring of dental fluorosis, sources of fluoride), interpretation and comparison of findings are difficult. Most of the studies in category (i) seem to indicate that the first year of life is a crucial period for the development of dental fluorosis. Pendrys et al. (7, 8), however, found that teeth exposed to fluoride supplements throughout the 3rd to the 6th year of life were equally at risk. Findings in category (ii) are more ambiguous, but a majority of reports support findings from animal studies stating that teeth in the transitional and early maturation stages (i.e. later stages of mineralization) rather than teeth in the secretory (early) stage are particularly susceptible

Table 1. The literature search strategy—MedLine\*

| No. | Records | Request                |
|-----|---------|------------------------|
| 1   | 1,537   | Fluorosis              |
| 2   | 215,462 | Dental                 |
| 3   | 1,178   | Fluorosis dental       |
| 4   | 116,852 | Risk factors           |
| 5   | 3,941   | Risk assessment        |
| 6   | 9,146   | Disease susceptibility |
| 7   | 6,387   | Dentition              |
| 8   | 39,210  | Permanent              |
| 9   | 705     | Dentition permanent    |
| 10  | 128,637 | #4 or #5 or #6         |
| 11  | 2       | #3 and #10 and #9      |
| 12  | 45      | #3 and #10             |
| 13  | 311,912 | Risk                   |
| 14  | 57,166  | Susceptible**          |
| 15  | 362,964 | #13 or #14             |
| 16  | 109     | #3 and #15             |
| 17  | 109     | #16 or #12 or #11      |

\* The same search strategy was used on the ISI database, and 69 studies were identified.

\*\* Truncated request.

(13–15). Ishii and Suckling (16) found two ‘at-risk periods’ for the development of moderate to severe fluorosis in the incisors; (i) starting at birth and ending relatively early in the developmental period of the tooth, and (ii) starting later during the mineralization period and ending shortly before eruption.

The lack of agreement between results from these two study designs may be due to the fact that dental fluorosis mirrors not only the daily fluoride load, but also the duration of fluoride exposure. Thus, an early introduction, e.g. to high-fluoride water, in most cases also indicates a long period of exposure. For category (ii), a reduction in daily fluoride intake in children while the tooth is in the maturation phase of development reflects a longer duration of fluoride exposure as compared to children experiencing fluoride reduction while the teeth are in the secretory phase of enamel formation. One problem encountered in evaluating the studies was inherent differences in study design and lack of statistical power due to small sample sizes.

The present review of the literature was done in order to up-date present knowledge, especially regarding the existence or non-existence of special ‘risk periods’. To avoid bias and confirm or refute individual findings, so-called meta-analytical methods have been developed to aggregate information from different studies and provide quantitative estimates of the effect. In the present study, data from relevant literature were combined statistically to provide a quantitative estimate of the strength of association between the period of exposure and risk of disease.

Hence, the purpose of this paper is threefold: (i) to present a systematic review of the existing dental literature dealing with periods of susceptibility to fluorosis in incisors, (ii) to assess the credibility of the various ‘risk period’

findings, and, finally, (iii) to test the hypothesis that these studies may measure *duration* of fluoride exposure rather than identify critical periods for developing dental fluorosis.

## Material and methods

### Literature search

A computerized literature search was made to identify relevant studies. A professional librarian specializing in literature search in medical topics carried out a search in the MedLine (WinSPIRS 2.0) and ISI databases (searches carried out May 8th, 1998). The strategy is given in *Table 1*. The databases cover the literature since 1966 (MedLine) and 1992 (ISI database). A classical study published in 1933 was also included (11).

A systematic search for unpublished studies was not carried out, but one manuscript was obtained and included in the meta-analysis (17). Authors of 2 papers were contacted in order to get information that was not easily deducible from their papers. However, a reply was obtained from the authors of one paper only (18).

### Inclusion criteria

To be included in the meta-analysis, the following requirements had to be met: The study had to assess dental fluorosis according to periods of fluoride exposure for the incisors, either directly (hypothesis testing) or indirectly (hypothesis forming). The studies included were classified as follows: Group (i), studies in individuals who had been introduced to fluoride at different ages during the enamel formation, and Group (ii), studies in children who, after having been exposed to fluoride from birth experienced an abrupt reduction in daily fluoride intake at a given time during the first 4 years of life, e.g. when a high-fluoride water source was replaced by low-fluoride water.

Furthermore, the number of cases versus controls (fluorosis versus non-fluorosis) for the incisors had to be given, as well as the age when first exposed to fluoride (or removed from the relevant fluoride source), or this had to be deduced with certainty. Alternatively, papers had to give the odds ratio (OR), standard error (SE) and the 95% confidence interval (CI) for the different groups. All indices of dental fluorosis were allowed, as the dependent variable was the prevalence of dental fluorosis and not the severity. If one index was applied in several of the included studies, a separate analysis was carried out based on these studies. There was no restriction on fluoride sources. A separate analysis for each fluoride source was carried out if the number of included studies allowed this.

Group (i) reports were divided into: (1) exposure to fluoride during the first 2 years of life and (2) exposure to fluoride only after the 2nd year of life (*Table 2*). Because some studies made the distinction between “early

Table 2. Studies with the Group (i) design included in the meta-analysis (in chronological order)

| Study no. | Authors (ref. no.)        | Year | Index used for scoring dental fluorosis | Fluoride source* | N   | Introduced to fluoride (age in months) | OR     | SE   | 95% CI                  |
|-----------|---------------------------|------|---|------------------|-----|--|--------|------|-------------------------|
| 1         | Forsman (20) **           | 1974 | Dean                                    | 1                | 7   | Early (0–24);<br>Late (>24)            | 0.0001 | 116  | 0.00–1.10 <sup>95</sup> |
| 2         | Holm & Andersson (6)      | 1982 | TFI                                     | 2, 3             | 86  | Early (6–23);<br>Late (≥24)            | 3.38   | 0.61 | 1.02–11.19              |
| 3         | Osuji et al. (21)         | 1988 | TFI                                     | (1), 2           | 139 | Early (0–24);<br>Late (>24)            | 11.92  | 0.41 | 5.31–26.80              |
| 4         | Lalumandier & Rozier (22) | 1995 | TSFI                                    | 2                | 113 | Early (0–23);<br>Late (≥24)            | 3.05   | 0.50 | 1.17–8.01               |
| 5         | Ismail & Messer (9)       | 1996 | FRI                                     | 1                | 48  | Early (0–23);<br>Late (≥24)            | 14.87  | 0.72 | 3.60–61.37              |
| 6         | Wang et al. (18)          | 1997 | TFI                                     | 2, 3             | 253 | Early (0–20);<br>Late (≥21)            | 7.22   | 0.56 | 2.40–21.70              |
| 7         | Bårdsen & Bjorvatn (17)   | 1998 | TFI                                     | 1 (2, 3)         | 66  | Early (0–24);<br>Late (>24)            | 11.32  | 0.62 | 3.39–37.87              |

\* 1: water, 2: fluoridated toothpaste, 3: fluoride supplements.

\*\* The study contains an empty (zero) cell in the 2 × 2 table.

exposure” and “later exposure” at for example 23 months of age and some used the 24th month for the distinction, this dichotomizing will not be exhaustive and exclusive. For Group (ii) reports, the fluoride exposure was divided into: (1) reduction of fluoride intake early in life (exposure to fluoride only during the first 2 years of life) and (2) reduction in fluoride intake later in life (exposure to fluoride for more than 2 years out of the first 4 years of life) (Table 3).

As both types of reports may assess duration of fluoride exposure rather than critical periods for developing dental fluorosis in the incisors, alternative analyses were carried out. The grouping was as follows: (a) long exposure (exposed to fluoride for more than 2 of the first 4 years of life), (b) short exposure (exposed less than 2 of the first 4 years of life) (Table 4).

#### Analysing the data, meta-analysis

Results from each of the included studies were entered into a data matrix (SPSS-PC, Version 7.5). Logistic regression analysis was used to calculate the OR, the 95% CI and the SE of the OR for each study (depen-

dent variable: fluorosis/not fluorosis, independent variables: groups according to age at start or age at end of fluoride exposure, as well as duration of fluoride exposure—weighted by the number of participants in each study).

If the data were homogeneous with respect to study features, the fixed effect model was applied to calculate the overall effect size (summary OR), as described by Fleiss (19). With lack of homogeneity, the random effects model was used to calculate the summary OR (19).

## Results

### Literature search

One-hundred-and-nine papers were identified through the MedLine search back to 1966, and 69 through the search in the ISI database. The findings overlapped in 37 cases, giving a total of 141 relevant papers. One unpublished paper (in manuscript) was obtained and included in the meta-analysis (17). Furthermore, one paper published before 1966 was included (11), giving a total of

Table 3. Studies with the Group (ii) design included in this meta-analysis (in chronological order)

| Study no. | Authors (ref. no.)       | Year | Index used for scoring dental fluorosis | Fluoride source* | N   | Reduction in fluoride intake (age in months) | OR    | SE    | 95% CI                 |
|-----------|--------------------------|------|---|------------------|-----|--|-------|-------|------------------------|
| 8         | McKay (11)               | 1933 | —                                       | 1                | 43  | Early (0–24)<br>Late (>24)                   | 10.12 | 0.91  | 1.70–60.2              |
| 9         | Ishii & Suckling (12) ** | 1986 | Dean                                    | 1 (2)            | 16  | Early (0–24)<br>Late (>24)                   | 36670 | 85.64 | 0.0–2.92 <sup>77</sup> |
| 10        | Evans & Stamm (10)       | 1991 | Dean                                    | 1 (2)            | 940 | Early (0–24)<br>Late (>24)                   | 1.75  | 0.17  | 1.25–2.47              |

\* 1: water, 2: fluoridated toothpaste, 3: fluoride supplements.

\*\* The study contains an empty (zero) cell in the 2 × 2 table.

Table 4. The pool of Group (i) and (ii) design included in the meta-analysis controlling for duration of exposure

| Study no. | Authors (ref. no.)        | Year | Index used for scoring dental fluorosis | Fluoride source* | N   | Duration of fluoride exposure (months)** | OR     | SE    | 95% CI                  |
|-----------|---------------------------|------|---|------------------|-----|--|--------|-------|-------------------------|
| 1         | Forsman (20)***           | 1974 | Dean                                    | 1                | 7   | Long (>23);<br>Short (≤23)               | 0.0001 | 116   | 0.00–1.10 <sup>95</sup> |
| 2         | Holm & Andersson (6)      | 1982 | TFI                                     | 2, 3             | 86  | Long (>24);<br>Short (≤24)               | 3.38   | 0.61  | 1.02–11.19              |
| 3         | Osuji et al. (21)         | 1988 | TFI                                     | (1), 2           | 139 | Long (>23);<br>Short (≤23)               | 11.92  | 0.41  | 5.31–26.80              |
| 4         | Lalumandier & Rozier (22) | 1995 | TSEFI                                   | 2                | 113 | Long (>24);<br>Short (≤24)               | 3.05   | 0.50  | 1.17–8.01               |
| 5         | Ismail & Messer (9)       | 1996 | FRI                                     | 1                | 48  | Long (>24);<br>Short (≤24)               | 14.87  | 0.72  | 3.60–61.37              |
| 6         | Wang et al. (18)          | 1997 | TFI                                     | 2                | 253 | Long (>27);<br>Short (≤27)               | 7.22   | 0.56  | 2.40–21.70              |
| 7         | Bårdsen & Bjorvatn (17)   | 1998 | TFI                                     | 1, (2, 3)        | 66  | Long (>23);<br>Short (≤23)               | 11.32  | 0.62  | 3.39–37.87              |
| 8         | McKay (11)                | 1933 | –                                       | 1                | 43  | Long (>23);<br>Short (≤23)               | 10.12  | 0.91  | 1.70–60.20              |
| 9         | Ishii & Suckling (12)***  | 1986 | Dean                                    | 1                | 16  | Long (>23);<br>Short (≤23)               | 36670  | 85.64 | 0.0–2.92 <sup>77</sup>  |
| 10        | Evans & Stamm (10)        | 1991 | Dean                                    | 1                | 940 | Long (>23);<br>Short (≤23)               | 1.75   | 0.17  | 1.25–2.47               |

\* 1: water, 2: fluoridated toothpaste, 3: fluoride supplements.

\*\* Duration of exposure, out of the first four years.

\*\*\* The study contains an empty (zero) cell in the 2 × 2 table.

143 papers. Only 10 of the studies satisfied the criteria for inclusion in the meta-analysis; 7 were Group (i) (6,

9, 17, 18, 20–22) and 3 Group (ii) (10–12) studies (Tables 2 and 3, respectively). The excluded studies, and the main reason for exclusion, are listed in Table 5.

Table 5. Studies excluded and main reason for rejection

| Main reason for exclusion, based on the topic described in the paper  | References    |
|---|---------------|
| Reporting on fluorosis, but not reporting age when introduced to fluoride (no control as to lifelong residency in high-fluoridated areas) | (8, 23–44)    |
| Scores for dental fluorosis not recorded/specified in the maxillary central incisors  | (5, 45–49)    |
| Age groups not suitable or overlapping groups   | (7, 50–52)    |
| Overview/review papers or safety considerations/guidelines  | (53–81)       |
| Histology/histopathology/morphology of fluorosed enamel   | (82–86)       |
| Reporting on dietary fluoride or fluoride concentrations in prophylactic fluoride products and/or the use of these                        | (37, 87–106)  |
| Including only enamel defects in deciduous teeth  | (107)         |
| Reporting on enamel fluoride concentrations and/or dental caries  | (108–116)     |
| Trends in prevalence of dental fluorosis  | (117)         |
| Aesthetic concerns of dental fluorosis  | (118–120)     |
| Material already presented in another included paper  | (16, 121–123) |
| Environmental effects of fluoride and/or fluoride in waters   | (124–129)     |
| Describing index/indices/techniques for scoring of dental fluorosis   | (130–132)     |
| Non-English paper   | (133–139)     |
| Other reason  | (140–151)     |

#### *Studies including subjects introduced to fluoride at different ages during enamel formation*

Children who were introduced to fluoride during the first 2 years of life had a significantly higher risk of developing dental fluorosis compared to those exposed after only 2 years of age. The material was homogeneous with respect to the effect size and, hence, the fixed effect model was used to calculate OR (19). Compared to children exposed to fluoride only after 2 years of age, those who were introduced to fluoride during the first 2 years of life had an OR of 7.24 (95% CI; 4.71–11.13) for developing dental fluorosis (Table 6).

When the meta-analysis was repeated, including only the 4 studies using the TF Index, the OR for early introduction to fluoride was 8.41 (95% CI; 5.05–14.01) (Table 6). Furthermore, separate analyses were carried out for studies including children who were (1) exposed to water fluoride and (2) exposed to fluoride supplements and fluoridated toothpaste. The summary OR and the 95% CI for these calculations are given in Table 6. Finally, new calculations were carried out after one study (18) had been excluded due to different age classification (20 months of age versus >20 months of age), and another (20) because of one empty (zero) cell in the 2 × 2 table (which gives an

Table 6. The summary OR for the studies in Group (i) where the subjects were exposed to fluoride at different age; early introduction versus late introduction\*

| Studies (ref. no.)   | OR    | 95% CI     |
|--|-------|------------|
| All (6, 9, 17, 18, 20–22)  | 7.24  | 4.71–11.13 |
| All except the study by Wang et al.**<br>(6, 9, 17, 20–22)                           | 7.24  | 4.53–11.59 |
| All except the study by Forsman***<br>(6, 9, 17, 18, 21, 22)                         | 7.24  | 4.66–11.25 |
| All except the studies by Forsman and<br>Wang et al. (6, 9, 17, 21, 22)              | 7.24  | 4.53–11.59 |
| Studies using the TF Index (6, 17, 18, 21)   | 8.41  | 5.05–14.01 |
| Studies where the fluoride source was<br>water (9, 17, 20)                           | 12.26 | 6.69–22.65 |
| Studies where toothpaste and supplement<br>were the fluoride sources (6, 18, 21, 22) | 3.60  | 2.20–5.87  |

\* Fixed effect model.

\*\* Excluded in the calculation due to the deviant way of dichotomizing the age groups.

\*\*\* Excluded in the calculation due to the empty cell (zero) cell in the 2 × 2 table.

unreliable OR). The results show that the inclusion of these 2 studies did not influence the outcome (Table 6).

*Studies including subjects who, after having been exposed to fluoride from birth, experienced an abrupt reduction in daily fluoride intake at different ages during enamel formation*

The Group (ii) studies included subjects who were cut off from the fluoride source or had a substantial reduction in fluoride intake at a certain age (6 months to 6 years) (Table 3). The Group (ii) children showed a significant overall effect of age at which the reduction in daily fluoride intake took place. Compared to children who experienced a reduction in daily fluoride intake as early as during the first 2 years of life, the OR was 1.88 (95% CI; 1.35–2.61) for children to develop dental fluorosis if the fluoride intake was reduced only *after* 2 years of age (i.e. the children had been exposed for more than 2 of the first 4 years of life).

Only 3 papers reported Group (ii) studies, all with water as the fluoride vehicle. One of the studies (11) was carried out before any fluorosis indices had been established, and

Table 7. The summary OR for the studies in Group (ii) where the subjects were introduced to fluoride at birth and then protected from the source at different age; reduction in fluoride intake late versus early\*

| Studies (ref. no)                                      | OR   | 95% CI    |
|--|------|-----------|
| All (10–12)  | 1.88 | 1.35–2.61 |
| All except the study by Ishii & Suckling**<br>(10, 11) | 1.88 | 1.35–2.61 |

\* Fixed effect model.

\*\* Excluded in the calculation due to the empty (zero) cell in the 2 × 2 table.

findings are given as pure descriptions of the examined tooth surfaces. The other 2 studies used Dean's Index. Separate analyses for the different indices were not done, but an alternative calculation was carried out after excluding the study by Ishii & Suckling (9), which had one empty (zero) cell in the 2 × 2 table. The results are given in Table 7.

#### Duration of fluoride exposure

The meta-analysis using duration of exposure as the independent variable (*long exposure*: >2 years out of the first 4 years, as compared to *short exposure*: <2 years) showed that children with a history of long exposure had a significantly higher risk of developing dental fluorosis compared with children with a short exposure. The summary OR was 5.83 (95% CI; 2.83–11.94) (Table 8).

When the meta-analysis was repeated including only the studies using the *TF Index* for scoring dental fluorosis (all studies from Group (i)), the OR was 8.41 (95% CI; 5.05–14.01) (Table 8). Similar analyses were carried out for studies using *Dean's Index*, for studies involving subjects *exposed to water fluoride*, and for studies related to children *exposed to fluoride toothpaste and supplements* only. The OR varied from 1.51 to 5.87 depending on the number of studies included and indices used (Table 8). All ORs were statistically significant at the 5% level.

## Discussion

Meta-analyses can organize results and thereby facilitate new findings, or put old findings in a new perspective. However, they also raise problems. A frequent criticism is that the studies available for analysis will represent a *biased* sample of all relevant studies carried out (152). As pointed out by McNemar (153), published studies tend to have larger *effect size* than those remaining unpublished. Hence, the estimate of the effect size from meta-analysis will be biased upward. This kind of bias, however, applies also to the narrative review. In the present study, both the hypothesis-forming and the hypothesis-testing studies have been included, and it is postulated that the availability bias should thereby be reduced.

A meta-analysis based on several randomized controlled trials is, in fact, a systematic review providing *new* data of high quality. Case-control studies were the basis for this meta-analysis. Improvement in the design and analysis of case-control studies has recently improved the quality of these studies and thereby eliminated much of the criticism previously raised (154). A disease with a long latency period such as dental fluorosis is well suited for case-control studies (154).

For meta-analysis to function, well-defined criteria for the inclusion of studies are required. The selection of studies, then, will be based on strict distinctions ('ex-

Table 8. The summary OR for the duration of fluoride exposure; long exposure versus short exposure (Group (i) and (ii))

| Studies (ref. no.)  | Factor analysis | OR   | 95% CI     |
|---|-----------------|------|------------|
| All (6, 9–12, 17, 18, 20–22)  | Random effect   | 5.83 | 2.83–11.94 |
| All except the study by Wang et al.* (6, 9–12, 17, 20–22)                                 | Random effect   | 5.87 | 2.66–12.94 |
| All except the studies by Forsman and Ishii & Suckling** (6, 9–11, 17, 18, 21, 22)        | Random effect   | 5.87 | 2.83–12.18 |
| All except the studies by Wang et al., Forsman and Ishii & Suckling (6, 9–11, 17, 21, 22) | Random effect   | 5.70 | 2.59–12.55 |
| Studies using the TF Index (6, 17, 18, 21)  | Fixed effect    | 8.41 | 5.05–14.01 |
| Studies using the Dean's Index (10, 12, 20)   | Random effect   | 1.51 | 1.11–2.05  |
| Studies where the fluoride source was water (9–12, 17, 20)                                | Fixed effect    | 2.32 | 1.71–3.15  |
| Studies where the fluoride sources were toothpaste and supplement (6, 18, 21, 22)         | Fixed effect    | 3.60 | 2.20–5.87  |

\* Excluded in the calculation due to the deviant way of dichotomizing the age groups.

\*\* Excluded in the calculation due to the empty (zero) cell in the  $2 \times 2$  table.

haustive and exclusive'), such as age. However, in the present study, the use of certain indefinite categories was accepted in the grouping of subjects according to months of fluoride exposure (overlapping at class boundaries). As there will always be some variation in the timing of enamel formation, deviation of 1 month (23 or 24 months of age) is not expected to influence the outcome significantly.

Despite the attempts to achieve objectivity in reviewing scientific data, considerable subjective judgement is required, not the least when deciding whether and how to use meta-analysis. Thus, as the quality of included studies can vary, meta-analyses may utilize different principles of weighting studies—thereby again introducing another subjective component. To avoid further bias, the present analysis was carried out considering only the criteria for inclusion as given in the material and method section.

The discrepancy between the number of papers identified in the literature search (143) and those actually included in the meta-analysis (10) might indicate too *wide* a search design compared to criteria for inclusion. This was accepted, however, as *risk periods* is not a *MESH term* in the MedLine database and the possibility for overlooking relevant papers was great.

The biochemical mechanisms leading to dental fluorosis in humans need further elucidation (15, 79, 155). According to DenBesten & Thariani (155), both animal and human studies indicate that teeth are particularly susceptible to fluoride during the transition/early-maturation stage of enamel formation. However, high fluoride intake during the secretory stage may also increase the risk. A study by Larsen et al. (50) showed that children starting to take fluoride supplements after the age of 2.5 years might develop dental fluorosis, i.e. even during the maturation stage teeth must be sensitive to fluoride exposure. The present qualitative and quantitative review found that early exposure to fluoride gave a significantly higher risk for developing dental fluorosis as compared to fluoride exposure later during the period of tooth formation. This would seem to support the theory that teeth are most susceptible to dental fluorosis during the

secretory phase of enamel formation. The Group (ii) design of studies, however, indicated that the risk was high also in children who experienced the reduction in daily fluoride exposure later, i.e. during the transitional and maturation phase of enamel formation.

As demonstrated by Table 8, long-time exposure to fluoride during the time of tooth formation significantly increased the risk for developing dental fluorosis. However, fairly big differences exist in the effect size for the different analyses. The most marked difference in OR was found to be between studies where the TF Index was used and those where Dean's Index was used in the assessment of fluorosis (Table 8). This may be because the TF Index is more sensitive and specific than Dean's Index (3), hence the TF Index may give a higher number of cases with dental fluorosis than Dean's Index. As studies using the TF Index for scoring of dental fluorosis were found only in Group (i), the sensitivity of this index may explain some of the difference in OR values found between Groups (i) and (ii).

Fluoride is known to be absorbed by bone and may also easily be released from bone (156). Information related to the fluoride removal (back-exchange) after a reduction in daily fluoride intake is scarce. However, it has been indicated that fluoride-induced disturbances in the amelogenesis occurs in the absence of elevated plasma fluoride concentrations in rat (157, 158). If the skeleton acts as an internal fluoride source after cessation of fluoride intake, this may interfere with the findings—especially for the analysis based on the Group (ii) studies—and add to the difference in the summary OR between Groups I and II.

The literature discussing risk periods related to dental fluorosis is scarce and different study designs make comparison and interpretation difficult. However, the results from the two different study designs included in the present analysis are compatible when the duration of fluoride exposure is assessed: Long exposure time (more than 2 of the first 4 years) increased the risk for developing dental fluorosis in the maxillary permanent incisors. On the other hand, based on the available epidemiological studies in humans, neither the secretory phase nor the

maturation phase of enamel formation can be singled out as being *the* high-risk period for the development of dental fluorosis in maxillary permanent central incisors.

*Acknowledgements.*—Sincere thanks to Dr Amid I. Ismail, who willingly gave important and encouraging information to initiate this meta-analysis. Advice in statistical analysis from Dr Håkon Gjessing is highly appreciated, and I am grateful to Svønn Sivertsen for carrying out the literature search. I am also indebted to Dr Kjell Björvatn and Dr Ola Haugejorden for invaluable criticism and advice during the preparation of the manuscript.

## References

- Dean HT, Arnold FA, Elvove E. Domestic water and dental caries. V. Additional studies of the relation of fluoride domestic water to dental caries experience in 4,425 white children aged 12–14 years in 13 cities in 4 states. *Public Health Rep* 1942;57:1155–79.
- Dean HT. *Epidemiological Studies in the United States*. In: Moulton FR, editor. *Dental caries and fluorine*. Washington D.C.: American Association for the Advancement of Science; 1946. p. 5–31.
- Thylstrup A, Fejerskov O. Clinical appearance of dental fluorosis in permanent teeth in relation to histologic changes. *Community Dent Oral Epidemiol* 1978;6:315–28.
- Nielsen HG, Ravn JJ. A radiographic study of mineralization of permanent teeth in a group of children aged 3–7 years. *Scand J Dent Res* 1976;84:109–18.
- Aasenden R, Peebles TC. Effects of fluoride supplementation from birth on human deciduous and permanent teeth. *Arch Oral Biol* 1974;19:321–6.
- Holm AK, Andersson R. Enamel mineralization disturbances in 12-year-old children with known early exposure to fluorides. *Community Dent Oral Epidemiol* 1982;10:335–9.
- Pendrys DG, Katz RV. Risk of enamel fluorosis associated with fluoride supplementation, infant formula, and fluoride dentifrice use. *Am J Epidemiol* 1989;130:1199–208.
- Pendrys DG, Katz RV, Morse DE. Risk factors for enamel fluorosis in a fluoridated population. *Am J Epidemiol*, 1994; 140:461–71.
- Ismail AI, Messer JG. The risk of fluorosis in students exposed to a higher than optimal concentration of fluoride in well water. *J Publ Health Dent* 1996;56:22–7.
- Evans WR, Stamm JW. An epidemiologic estimate of the critical period during which human maxillary central incisors are most susceptible to fluorosis. *J Publ Health Dent* 1991; 51:251–9.
- McKay FS. Mottled enamel: the prevention of its further production through a change of the water supply at Oakley, IDA. *J Am Dent Assoc* 1933;20:1137–49.
- Ishii T, Suckling G. The appearance of tooth enamel in children ingesting water with a high fluoride content for a limited period during early tooth development. *J Dent Res* 1986;65:974–7.
- Kierdorf U, Kierdorf H, Fejerskov O. Fluoride-induced developmental changes in enamel and dentine of European roe deer (*Capreolus capreolus* L.) as a result of environmental pollution. *Arch Oral Biol* 1993;38:1071–81.
- Kierdorf U, Kierdorf H, Sedlacek F, Fejerskov O. Structural changes in fluorosed dental enamel of red deer (*Cervus elaphus* L.) from a region with severe environmental pollution by fluorides. *J Anat* 1996;188:183–95.
- Richards A, Kragstrup J, Josephsen K, Fejerskov O. Dental fluorosis developed in post secretory enamel. *J Dent Res* 1986;65:1406–9.
- Ishii T, Suckling G. The severity of dental fluorosis in children exposed to water with a high fluoride content for various periods of time. *J Dent Res* 1991;70:952–6.
- Bårdsen A, Björvatn K. Risk of dental fluorosis when exposed to high-fluoride water at different age. *J Dent Res* 1998;77 (Special Issue B, Abstracts of Papers): 700.
- Wang NJ, Gropen AM, Øgaard B. Risk factors associated with fluorosis in a non-fluoridated population in Norway. *Community Dent Oral Epidemiol* 1997;25:396–401.
- Fleiss JL. The statistical basis of meta-analysis. *Statistical Meth Med Res* 1993;2:121–145.
- Forsman B. Dental fluorosis and caries in high-fluoride districts in Sweden. *Community Dent Oral Epidemiol* 1974;2:132–48.
- Osuji O, Leake JL, Chipman ML, Nikiforuk G, Locker D, Levine N. Risk factors for dental fluorosis in a fluoridated community. *J Dent Res* 1988;67:1488–92.
- Lalumandier JA, Rozier RG. The prevalence and risk factors of fluorosis among patients in a pediatric dental practice. *Pediatr Dent* 1995;17:19–25.
- Hallett GE, Porteous JR. Observations on the dental health of young adults in Hartlepool and Newcastle. *Br Dent J* 1970; 128:171–4.
- Keene HJ, Grossman FD, Pederson ED, Mellberg JR, Nicholson CR. Caries experience and fluoride concentrations of surface enamel in dental fluorosis patients. *Caries Res* 1975;9:247–50.
- Rozier RG, Dudley GG. Dental fluorosis in children exposed to multiple sources of fluoride: implications for school fluoridation programs. *Public Health Rep* 1981;96:542–6.
- Butler WJ, Segreto V, Collins E. Prevalence of dental mottling in school-aged lifetime residents of 16 Texas communities. *Am J Public Health* 1985;75:1408–12.
- Granath L, Widenheim J, Birkhed D. Diagnosis of mild enamel fluorosis in permanent maxillary incisors using two scoring systems. *Community Dent Oral Epidemiol* 1985;13:273–6.
- Manji F, Baelum V, Fejerskov O, Gemert W. Enamel changes in two low-fluoride areas of Kenya. *Caries Res* 1986;20:371–80.
- Eklund SA, Burt BA, Ismail AI, Calderone JJ. High-fluoride drinking water, fluorosis and dental caries in adults. *J Am Dent Assoc* 1987;114:324–8.
- Wöltgens JHM, Etty EJ, Nieuwland WMD, Lyaruu DM. Use of fluorosis by young children and prevalence of mottled enamel. *Adv Dent Res* 1989;3:177–82.
- Ismail AI, Brodeur JM, Kavanagh M, Boisclair G, Tessier C, Picotte L. Prevalence of dental caries and dental fluorosis in students, 11–17 years of age, in fluoridated and non-fluoridated cities in Quebec. *Caries Res* 1990;24:290–7.
- Riordan PJ, Banks JA. Dental fluorosis and fluoride exposure in Western Australia. *J Dent Res* 1991;70:1022–8.
- Ismail AI, Shoveller J, Langille D, MacInnis W, McNally M. Should the drinking water of Truro, Nova Scotia, be fluoridated? Water fluoridation in the 1990's. *Community Dent Oral Epidemiol* 1993;21:118–25.
- Riordan PJ. Dental fluorosis, dental caries and fluoride exposure among 7-year-olds. *Caries Res* 1993;27:71–7.
- Clark DC, Hann HJ, Williamson MF, Berkowitz J. Influence of exposure to various fluoride technologies on the prevalence of dental fluorosis. *Community Dent Oral Epidemiol* 1994; 22:461–4.
- Awad MA, Hargreaves JA, Thompson GW. Dental caries and fluorosis in 7–9 and 11–14 year old children who received fluoride supplements from birth. *J Can Dent Assoc* 1994;60: 318–22.
- Ellwood RP, O'Mullane DM. Association between dental enamel opacities and dental caries in a North Wales population. *Caries Res* 1994;28:383–7.
- Grimaldo M, Botja Aburto VH, Ramirez AL, Ponce M, Diaz Barriga F. Endemic fluorosis in San Luis Potosi, Mexico. I. Identification of risk factors associated with human exposure to fluoride. *Environ Res* 1995;68:25–30.
- Skotowski MC, Hunt RJ, Levy SM. Risk factors for dental

- fluorosis in pediatric dental patients. *J Public Health Dent* 1995;55:154–9.
40. Wang LF, Huang JZ. Outline of control practice of endemic fluorosis in China. *Soc Sci Med* 1995;41:1191–5.
  41. Pendrys DG. Risk of fluorosis in a fluoridated population. Implication for the dentist and hygienist. *Am J Dent Assoc* 1995;126:1617–24.
  42. DiazBarriga F, Leyva R, Quistian J, Loyola Rodriguez JP, Pozos A, Grimaldo M. Endemic fluorosis in San Luis Potosi, Mexico. 4. Sources of fluoride exposure. *Fluoride* 1997;30:219–22.
  43. Heller KE, Eklund SA, Burt BA. Dental caries and dental fluorosis at varying fluoride concentrations. *J Public Health Dent* 1997;57:136–43.
  44. Clark DC, Berkowitz J. The influence of various fluoride exposures on the prevalence of esthetic problems resulting from dental fluorosis. *J Public Health Dent* 1997;57:144–9.
  45. Aasenden R, Peebles TC. Effects of fluoride supplementation from birth on dental caries and fluorosis in teenaged children. *Arch Oral Biol* 1978;23:111–5.
  46. Bagramian RA, Narendran S, Ward M. Relationship of dental caries and fluorosis to fluoride supplement history in a non-fluoridated sample of schoolchildren. *Adv Dent Res* 1989;3:161–7.
  47. Kumar JV, Green EL, Wallace W, Carnahan T. Trends in dental fluorosis and dental caries prevalences in Newburgh and Kingston, NY. *Am J Public Health* 1989;79:565–9.
  48. Somvanshi PR, Chaubey BS, Phadake RV, Sakharde PM. Fluorosis in Maharashtra. *J Assoc Physicians India* 1990;38:217–9.
  49. Bårdsen A, Klock KS, Bjorvatn K. Dental fluorosis in selected areas of Norway. Risk factors. *J Dent Res* 1997;76:1107.
  50. Larsen MJ, Richards A, Fejerskov O. Development of dental fluorosis according to age at start of fluoride administration. *Caries Res* 1985;19:519–27.
  51. Holt RD, Morris CE, Winter GB, Downer MC. Enamel opacities and dental caries in children who used a low fluoride toothpaste between 2 and 5 years of age. *Int Dent J* 1994;44:331–41.
  52. Pendrys DG, Katz RV, Morse DE. Risk factors for enamel fluorosis in a nonfluoridated population. *Am J Epidemiol* 1996;143:808–15.
  53. Heifetz SB, Horowitz HS. The amount of fluoride in current fluoride therapies: safety considerations for children. *ASDC J Dent Child* 1984;51:257–69.
  54. Ripa LW. Topical fluorides: a discussion of risks and benefits. *J Dent Res* 1987;66:1079–83.
  55. Riordan PJ. Guidelines for the use of dietary fluoride supplements in Australia. *Aust Dent J* 1989;34:359–62.
  56. Horowitz HS. Fluoride and enamel defects. *Adv Dent Res* 1989;3:143–6.
  57. Kaminsky LS, Mahoney C, Leach J, Melius J, Miller MJ. Fluoride: benefits and risks of exposure. *Crit Rev Oral Biol Med* 1990;1:261–81.
  58. Pendrys DG, Stamm JW. Relationship of total fluoride intake to beneficial effects and enamel fluorosis. *J Dent Res* 1990;69 (Spec No):529–38.
  59. Rugg-Gunn AJ, Murray JJ. Current issues in the use of fluorides in dentistry. *Dent Update* 1990;17:154–8.
  60. Ripa LW. A critique of topical fluoride methods (dentifrices, mouthrinses, operator-, and self-applied gels) in an area of decreased caries and increased fluorosis prevalence. *J Public Health Dent* 1991;51:23–41.
  61. From the Centers for Disease Control. Public Health Service report on fluoride benefits and risks. *JAMA* 1991;266:1061–2, 1066–7.
  62. Public Health Service report on fluoride benefits and risks *MMWR Morb Mortal Wkly Rep* 1991;40:1–8.
  63. Pendrys DG. Dental fluorosis in perspective. *J Am Dent Assoc* 1991;122:63–6.
  64. Newbrun E. Current regulations and recommendations concerning water fluoridation, fluoride supplements, and topical fluoride agents. *J Dent Res* 1992;71:1255–65.
  65. Szpunar SM, Burt BA. Evaluation of appropriate use of dietary fluoride supplements in the US. *Community Dent Oral Epidemiol* 1992;20:148–54.
  66. D'Hoore W, Van Nieuwenhuysen JP. Benefits and risks of fluoride supplementation: caries prevention versus dental fluorosis. *Eur J Pediatr* 1992;151:613–6.
  67. Kaminsky LS, Mahoney MC, Miller MJ. Fluoride: benefits and risk of exposure. The preparation of a report. *J Am Coll Dent* 1992;59:4–7.
  68. Levy SM. A review of fluoride intake from fluoride dentifrice. *ASDC J Dent Child* 1993;60:115–24.
  69. Riordan PJ. Fluoride supplements in caries prevention: a literature review and proposal for a new dosage schedule. *J Public Health Dent* 1993;53:174–89.
  70. Li Y. Fluoride: safety issues. *J Indiana Dent Assoc* 1993;72:22–6.
  71. Johnston DW. Current status of professionally applied topical fluorides. *Community Dent Oral Epidemiol* 1994;22:159–63.
  72. Levy SM. Review of fluoride exposure and ingestion. *Community Dent Oral Epidemiol* 1994;22:173–80.
  73. Ismail AI. Fluoride supplements: current effectiveness, side effects, and recommendations. *Community Dent Oral Epidemiol* 1994;22:164–72.
  74. Stookey GK. Review of fluorosis risk of self-applied topical fluorides: dentifrices, mouth rinses and gels. *Community Dent Oral Epidemiol* 1994;22:181–6.
  75. Lewis DW, Banting DW. Water fluoridation—current effectiveness and dental fluorosis. *Community Dent Oral Epidemiol* 1994;22:153–8.
  76. Horowitz HS. Commentary on and recommendations for the proper use of fluoride. *J Public Health Dent* 1995;55:57–62.
  77. Ismail AI. What is the effective concentration of fluoride. *Community Dent Oral Epidemiol* 1995;23:246–51.
  78. Holt RD, Murray JJ. Developments in fluoride toothpaste—an overview. *Community Dent Health* 1997;14:4–10.
  79. Withford GM. Determinants and mechanisms of enamel fluorosis. In: Ciba Foundation Symposium 205. *Dental Enamel*. Chichester: John Wiley & Sons; 1997. p. 226–45.
  80. Holloway PJ, Ellwood RP. The prevalence, causes and cosmetic importance of dental fluorosis in the United Kingdom: a review. *Community Dent Health*, 1997;14:148–55.
  81. Fluoridation debate *Environ Health Perspect* 1997;105:1176.
  82. Gustavson G, Sundström B. Enamel: morphological considerations. *J Dent Res* 1975;54 (Spec Issue No. B):114–20.
  83. Withford GM, Allmann DW, Shahed AR. Topical fluorides: effects on physiological and biochemical processes. *J Dent Res* 1987;66:1072–8.
  84. DenBesten P. Dental fluorosis: its use as a biomarker. *Adv Dent Res* 1994;8:105–10.
  85. Fejerskov O, Larsen MJ, Richards A, Baelum V. Dental tissue effects of fluoride. *Adv Dent Res* 1994;8:15–31.
  86. Aoba T. The effect of fluoride on apatite structure and growth. *Crit Rev Oral Biol Med* 1997;8:136–53.
  87. Baume LJ, Vulliemoz JP. Dietary fluoride uptake into the enamel of caries-susceptible “yellow” permanent teeth and of caries-resistant permanent and primary teeth of Polynesians. *Arch Oral Biol* 1970;15:431–43.
  88. Heifetz SB, Horowitz HS. Amount of fluoride in self-administered dental products: safety considerations for children. *Pediatrics* 1986;77:876–82.
  89. Pendrys DG, Morse DE. Use of fluoride supplementation by children living in fluoridated communities. *ASDC J Dent Child* 1990;57:343–7.
  90. Horowitz HS. The need for toothpaste with lower than conventional fluoride concentrations for preschool-aged children. *J Public Health Dent* 1992;52:216–21.
  91. Levy SM, Maurice TJ, Jakobsen JR. A pilot study of

- preschoolers' use of regular-flavored dentifrices and those flavored for children. *Pediatr Dent* 1992;14:388-91.
92. Levy SM, Maurice TJ, Jakobsen JR. Feeding patterns, water sources and fluoride exposure of infants and 1 year olds. *J Am Dent Assoc* 1993;124:65-9.
  93. Levy SM, Maurice TJ, Jakobsen JR. Dentifrice use among preschool children. *J Am Dent Assoc* 1993;124:57-60.
  94. Mwaniki DL, Opinya GN. Fluoride and abrasive content in commonly used dentifrices on the Kenyan market. *East Afr Med J* 1993;70:288-90.
  95. Nourjah P, Horowitz AM, Wagener DK. Factors associated with the use of fluoride supplements and fluoride dentifrice by infants and toddlers. *J Public Health Dent* 1994;54:47-54.
  96. Bergmann KE, Bergmann RL. Salt fluoridation and general health. *Adv Dent Res* 1995;9:138-43.
  97. Levy SM, Kohout FJ, Guha Chowdhury N, Kiritsy MC, Heilman J, Wefel JS. Infants' fluoride intake from drinking water alone, and from water added to formula, beverages, and food. *J Dent Res* 1995;74:1399-407.
  98. Levy SM, Kiritsy MC, Warren J. Sources of fluoride intake in children. *J Public Health Dent* 1995;55:39-52.
  99. Van Winkle S, Levy SM, Kiritsy MC, Heilman JR, Wefel JS, Marshall T. Water and formula fluoride concentrations: significance for infants fed formula. *Pediatr Dent* 1995;17:305-10.
  100. Mwaniki DL, Gikunju JK. Fluoride concentrations in tissue of fish from low fluoride fresh water lakes in Kenya. *Discovery and Innovation* 1995;7:173-6.
  101. Mascarenhas AK, Burt BA. Fluoride toothpaste. A risk factor in dental fluorosis. *J Dent Res* 1996;75:1361.
  102. Silva M, Reynold EC. Fluoride content of infant formulae in Australia. *Aust Dent J* 1996;41:37-42.
  103. Levy SM, Kiritsy MC, Slager SL, Warren J, Kohout FJ. Patterns of fluoride dentifrice use among infants. *Pediatr Dent* 1997;19:50-5.
  104. Adair SM, Piscitelli WP, McKnight HC. Comparison of the use of a child and a adult dentifrice by a sample of preschool children. *Pediatr Dent* 1997;19:99-103.
  105. Heilman JR, Kiritsy MC, Levy SM, Wefel JS. Fluoride concentrations of infant foods. *J Am Dent Assoc* 1997;128:857-63.
  106. Neesham DC. Fluoride concentration in AgF and dental fluorosis. *Aust Dent J* 1997;42:268-9.
  107. Milsom KM, Woodward M, Haran D, Lennon MA. Enamel defects in the deciduous dentition as a potential predictor of the defects in the permanent dentition of 8- and 9-year-old children in fluoridated Cheshire, England. *J Dent Res* 1996;75:1015-8.
  108. Keene HJ, Melberg JR, Nicholson CR. History of fluoride, dental fluorosis and concentrations of fluoride in surface layer of enamel of caries-free naval recruits. *J Public Health Dent* 1973;33:142-8.
  109. Larson RH. Animal studies relating to caries inhibition by fluoride. *Caries Res* 1977;11 Suppl 1:42-58.
  110. Leverett DH, Featherstone JD, Proskin HM, Adair SM, Eisenberg A, Mundorff Shrestha SA, et al. Caries risk assessment by a cross-sectional discriminant model. *J Dent Res* 1993;72:529-37.
  111. Leverett DH, Proskin HM, Featherstone JD, Adair SM, Eisenberg AD, Mundorff Shrestha SA, et al. Caries risk assessment in a longitudinal discrimination study. *J Dent Res* 1993;72:538-43.
  112. Lewis DW, Ismail AI, Goldblom R, Battista RN, Anderson G, Beaulieu McLeod R, et al. Periodic health examination, 1995 update. 2. Prevention of dental caries. *Can Med Assoc J* 1995;152:836-46.
  113. Brown IJ, Selwitz RH. The impact of recent changes in the epidemiology of dental caries on guidelines for the use of dental sealants. *J Public Health Dent* 1995;55:274-91.
  114. Ellwood RP, O'Mullane D. The association between developmental enamel defects and caries in populations with and without fluoride in their drinking water. *J Public Health Dent* 1996;56:76-80.
  115. Riordan PJ. The place of fluoride supplements in caries prevention today. *Aust Dent J* 1996;41:335-42.
  116. Leverett DH, Adair SM, Vaughan BW, Proskin HM, Moss ME. Randomized clinical trials of the effect of prenatal fluoride supplements in preventing dental caries. *Caries Res* 1997;31:174-9.
  117. Clark DC. Trends in prevalence of dental fluorosis in North-America. *Community Dent Oral Epidemiol* 1994;22:148-52.
  118. Clark DC, Hann HJ, Williamson MF, Berkowitz J. Aesthetic concerns of children and parents in relation to different classification of the Tooth Surface Index of Fluorosis. *Community Dent Oral Epidemiol* 1993;21:360-4.
  119. Riordan PJ. Perception of dental fluorosis. *J Dent Res* 1993;72:1268-74.
  120. Hawley GM, Ellwood RP, Davies RM. Dental caries, fluorosis and the cosmetic implications of different TF scores in 14-year-old adolescents. *Community Dent Health* 1996;13:189-92.
  121. Evans RW. Changes in dental fluorosis following an adjustment to the fluoride concentration of Hong Kong's water supplies. *Adv Dent Res* 1989;3:154-60.
  122. Evans RW. An epidemiological assessment of the chronological distribution of dental fluorosis in human maxillary central incisors. *J Dent Res* 1993;72:883-90.
  123. Evans RW, Darvell BW. Refining the estimate of the critical period for susceptibility to enamel fluorosis in human maxillary central incisors. *J Public Health Dent* 1995;55:238-49.
  124. Marier JR. Some current aspects of environmental fluoride. *Sci Total Environ* 1977;8:253-65.
  125. Gaciri SJ, Davies TC. The occurrence and geochemistry of fluoride in some natural waters of Kenya. *J Hydrology* 1993;143:395-412.
  126. Sirikanth R, Khanam A, Rao AMM. Fluoride in borehole water in selected villages of Medak district, Andhra-Pradesh, India. *Fluoride* 1994;27:93-6.
  127. Toumba KJ, Levy S, Curzon ME. The fluoride content of bottled drinking waters. *Br Dent J* 1994;176:266-8.
  128. Rozier RG. A new area for community water fluoridation? Achievements after one-half century and challenges ahead. *J Public Health Dent* 1995;55:3-5.
  129. Spencer AJ, Slade GD, Davies M. Water fluoridation in Australia. *Community Dent Health* 1996;13 Suppl 2:27-37.
  130. Pendrys DG. The fluorosis risk index: a method for investigating risk factors. *J Public Health Dent* 1990;50:291-8.
  131. Rozier RG. Epidemiologic indices for measuring the clinical manifestations of dental fluorosis: overview and critique. *Adv Dent Res* 1994;8:39-55.
  132. Kingman A. Current techniques for measuring dental fluorosis: issues in data analysis. *Adv Dent Res* 1994;8:56-65.
  133. Kerebel B, Dubois T, Clergeau Gueritault S, Siau T. (The human dental enamel in endemic fluorosis). *SSO Schweiz Monatsschr Zahnheilkd* 1973;83:326-43.
  134. Brabant H, Klees L, Newesely H. (The environmental factor in the etiology and origin of dental caries). *Acta Stomatol Belg* 1974;71:169-210.
  135. Modica R, Benech A. (Dental caries and experimental fluorosis). *Minerva Stomatol* 1983;32:377-80.
  136. Sergio Gomez S, Weber A, Torres C. (Fluoride content of tea and amount ingested by children). *Odontol Child* 1989;37:251-5.
  137. Løkken P, Valderhaug J. (Fluorosis due to fluoride in drinking water. A health risk even in Norway?). *Tidsskr Nor Laegeforen* 1991;111:1506-7.
  138. Van Nieuwenhuysen JP, D'Hoore W. (Dental caries, fluoride tablets and enamel opacities). *Arch Fr Pediatr* 1992;49:617-21.
  139. Valova GA, Polzik EV, Zinger VE, Shcherbakov SV. (Dermatoglyphic indices in assessing the predisposition to occupational fluorosis). *Tsitol Genet* 1994;28:56-9.

140. Colquhoun J. New evidence on fluoridation. *Soc Sci Med* 1984;19:1239–46.
141. de Muniz BR. Epidemiologic oral health survey of Argentine children. *Community Dent Oral Epidemiol* 1985;13:328–33.
142. Rozier RG, Beck JD. Epidemiology of oral disease. *Curr Opin Dent* 1991;1:308–15.
143. Holloway PJ. International dental public health. *Curr Opin Dent* 1991;1:348–56.
144. Morgan MV, Wright FA, Matram ZN, Sundoro E, Chesters RK. The oral health status and health behaviour of 12 and 15-year-old adolescents in Jakarta, Indonesia. *Community Dent Health* 1992;9:171–9.
145. Jackson R, Kelly S, Noblitt T, Zhang W, Dunipace A, Li Y, et al. The effect of fluoride therapy on blood-chemistry parameters in osteoporotic females. *Bone and Mineral* 1994;27:13–23.
146. Clark MM, Album MM, Lloyd RW. Preventive dentistry and the family physician. *Fam Physician* 1996;53:619–26, 631–2.
147. DenBesten P, Ko HS. Fluoride levels in whole saliva of preschool children after brushing with 0.25 g (pea-sized) as compared to 1.0 g (full-brush) of a fluoride dentifrice. *Pediatr Dent* 1996;18:277–80.
148. Woodward GL, Leake JL, Main PA. Oral health and family characteristics of children attending private or public dental clinics. *Community Dent Oral Epidemiol* 1996;24:253–9.
149. Grimaldo M, Turrubiartes F, Milan J, Pozos A, Alfaro C, DiazBarriga F. Endemic fluorosis in San Luis Potosi, Mexico. 3. Screening for fluoride exposure with a geographic information system. *Fluoride* 1997;30:33–40.
150. Kumar J, Swango P, Lininger L, Leske G, Green E, Haley V. Effect of case definition on the association between risk factors and dental fluorosis. *J Dent Res* 1997;76:3041.
151. Roberts MW, Keels MA, Sharp MC, Lewis JL. Fluoride supplements prescribing and dental referral patterns among academic pediatricians. *Pediatrics* 1998;101:E61–4.
152. Hunter JE, Schmidt FL. *Methods of meta-analysis: correcting error and bias in research findings*. Newbury Park, London, New Delhi: Sage Publications; 1990. p. 592.
153. McNemar Q. At random: sense and nonsense. *Am Psychol* 1960;15:295–300.
154. Hennekens CH, Buring JE. Case-control studies. In: Mayrent SL, editor. *Epidemiology in medicine*. Little, Brown and Company: Boston/Toronto; 1987. p. 132–52.
155. DenBesten PK, Thariani H. Biological mechanisms of fluorosis and level and timing of systemic exposure to fluoride with respect to fluorosis. *J Dent Res* 1992;71:1238–43.
156. Li J, Nakagaki H, Kato K, Tsuboi S, Kato S, Morita I, et al. Effect of stopping fluoride administration on the distribution profiles of fluoride in three different kinds of rat bones. *Calcif Tissue Int* 1995;56:292–6.
157. Angmar-Månsson B, Whitford GM. Single fluoride dose and enamel fluorosis in the rat. *Caries Res* 1985;19:145–52.
158. Angmar-Månsson B, Lindh U, Withford GM. Enamel and dentin fluoride levels and fluorosis following single fluoride doses: a nuclear microprobe study. *Caries Res* 1990;24:258–62.

---

Received for publication 8 March 1999

Accepted 9 August 1999